



## Persistent Mixed-Valence (TTF)(2) (+.) Dyad of a Chiral Bis(binaphthol)-tetrathiafulvalene (TTF) Derivative

Submitted by Emmanuel Lemoine on Thu, 02/06/2014 - 11:15

Titre	Persistent Mixed-Valence (TTF)(2) (+.) Dyad of a Chiral Bis(binaphthol)-tetrathiafulvalene (TTF) Derivative
Type de publication	Article de revue
Auteur	Saad, Ali [1], Barrière, Frédéric [2], Levillain, Eric [3], Vanthuyne, Nicolas [4], Jeannin, Olivier [5], Fourmigué, Marc [6]
Editeur	Wiley
Type	Article scientifique dans une revue à comité de lecture
Année	2010
Langue	Anglais
Numéro	27
Pagination	8020-8028
Volume	16
Titre de la revue	Chemistry - A European Journal
ISSN	0947-6539
Résumé en anglais	<p>Mixed-valence dyadic [(TTF)(2)](+.) (TTF=tetrathiafulvalene) species the elementary building blocks of organic conductors-are usually too weakly associated to be observed in solution, unless covalently bound in dimers or physically constrained into a cage structure. We demonstrate here that a novel chiral tetrathiafulvalene functionalised with two 1,1'-binaphthol units (1) is able to associate in solution into persistent mixed-valence [(TTF)(2)](+.) dyadic moieties through a stereospecific recognition pattern. This redox active molecule exhibits different electrochemical and spectroscopic responses, as enantiopure RR, SS or meso isomers, a rare example of a chiral system in which different diastereoisomers do not exhibit the same electrochemical features, with a selective formation of the mixed-valence species in the enantiopure (RR)-1 or (SS)-1 isomers only, whereas the meso form does not show this association ability. A rationale for the selective self-association of the RR and SS enantiomers upon oxidation is provided, based on the different molecular geometries and accessibility of the TTF core toward the formation of the mixed-valence species.</p>
URL de la notice	<a href="http://okina.univ-angers.fr/publications/ua2692">http://okina.univ-angers.fr/publications/ua2692</a> [7]
DOI	10.1002/chem.200902050 [8]

### Liens

- [1] [http://okina.univ-angers.fr/publications?f\[author\]=2972](http://okina.univ-angers.fr/publications?f[author]=2972)
- [2] [http://okina.univ-angers.fr/publications?f\[author\]=2973](http://okina.univ-angers.fr/publications?f[author]=2973)
- [3] <http://okina.univ-angers.fr/eric.levillain/publications>
- [4] [http://okina.univ-angers.fr/publications?f\[author\]=2943](http://okina.univ-angers.fr/publications?f[author]=2943)

- [5] [http://okina.univ-angers.fr/publications?f\[author\]=2974](http://okina.univ-angers.fr/publications?f[author]=2974)
- [6] [http://okina.univ-angers.fr/publications?f\[author\]=2678](http://okina.univ-angers.fr/publications?f[author]=2678)
- [7] <http://okina.univ-angers.fr/publications/ua2692>
- [8] <http://dx.doi.org/10.1002/chem.200902050>

Publié sur *Okina* (<http://okina.univ-angers.fr>)